

University of Groningen

Genome Sequences of 12 Spore-Forming Bacillus Species, Comprising *Bacillus coagulans*, *Bacillus licheniformis*, *Bacillus amyloliquefaciens*, *Bacillus sporothermodurans*, and *Bacillus vallismortis*, Isolated from Foods

Krawczyk, Antonina O.; de Jong, Anne; Holsappel, Siger; Eijlander, Robyn T.; van Heel, Auke; Berendsen, Erwin M; Wells-Bennik, Marjon H.J.; Kuipers, Oscar P.

Published in:
Genome Announcements

DOI:
[10.1128/genomeA.00103-16](https://doi.org/10.1128/genomeA.00103-16)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2016

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Krawczyk, A. O., de Jong, A., Holsappel, S., Eijlander, R. T., van Heel, A., Berendsen, E. M., Wells-Bennik, M. H. J., & Kuipers, O. P. (2016). Genome Sequences of 12 Spore-Forming Bacillus Species, Comprising *Bacillus coagulans*, *Bacillus licheniformis*, *Bacillus amyloliquefaciens*, *Bacillus sporothermodurans*, and *Bacillus vallismortis*, Isolated from Foods. *Genome Announcements*, 4(3), 1-2. [e00103-16].
<https://doi.org/10.1128/genomeA.00103-16>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Genome Sequences of 12 Spore-Forming *Bacillus* Species, Comprising *Bacillus coagulans*, *Bacillus licheniformis*, *Bacillus amyloliquefaciens*, *Bacillus sporothermodurans*, and *Bacillus vallismortis*, Isolated from Foods

Antonina O. Krawczyk,^{a,c} Anne de Jong,^{a,c} Siger Holsappel,^a Robyn T. Eijlander,^{a,c} Auke van Heel,^a Erwin M. Berendsen,^{a,b,c} Marjon H. J. Wells-Bennik,^{b,c} Oscar P. Kuipers^{a,c}

Molecular Genetics, University of Groningen, Groningen, The Netherlands^a; NIZO food research, Ede, The Netherlands^b; Top Institute Food and Nutrition (TIFN), Wageningen, The Netherlands^c

Here, we report the draft genomes of twelve isolates of five different *Bacillus* species, all spore-forming, Gram-positive bacteria.

Received 27 January 2016 Accepted 17 March 2016 Published 12 May 2016

Citation Krawczyk AO, de Jong A, Holsappel S, Eijlander RT, van Heel A, Berendsen EM, Wells-Bennik MHJ, Kuipers OP. 2016. Genome sequences of 12 spore-forming *Bacillus* species, comprising *Bacillus coagulans*, *Bacillus licheniformis*, *Bacillus amyloliquefaciens*, *Bacillus sporothermodurans*, and *Bacillus vallismortis*, isolated from foods. *Genome Announc* 4(3):e00103-16. doi:10.1128/genomeA.00103-16.

Copyright © 2016 Krawczyk et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Oscar P. Kuipers, o.p.kuipers@rug.nl.

Bacillus species have the ability to form endospores (spores). *Bacillus* spores are ubiquitously present in soil, and transmission to food products can take place (1). Spores are highly resistant to environmental stresses and can food processing conditions. Germination of spores followed by growth may result in food spoilage (2). Here, we report the draft genome sequences of twelve strains belonging to five different *Bacillus* species that were isolated from foods: Four strains of *Bacillus coagulans*, four strains of *Bacillus licheniformis*, two strains of *Bacillus amyloliquefaciens*, one strain of *Bacillus sporothermodurans*, and one strain of *Bacillus vallismortis*. Comparison of the sequenced genomes with those of *B. subtilis* may provide insight into variations in the sporulation and germination processes (3). Furthermore, genome mining can provide insight into the genomic potential of strains in relation to predicted phenotypic traits and their ability to produce toxins involved in food poisoning, such as lichenysin in *B. licheniformis* (4).

Twelve strains of different isolation sources (Table 1), were

grown overnight in 10 ml of brain heart infusion (BHI) broth (Difco) at 37°C. The overnight cultures were diluted 100-fold in fresh medium and incubated at 37°C until the culture reached an optical density (at 660 nm) of approximately 0.5, and cells were then harvested by centrifugation at 5000 rcf. DNA was isolated as described previously (5). The isolated DNA was sheared to 500-bp fragments in a Covaris (KBioscience) ultrasonication device for preparing the paired-end NEB NextGen library preparation kit. The prepared libraries were 101 bases paired-end sequenced on an Illumina HiSeq2000 by multiplexing 12 samples per flow cell. *De novo* paired-end assembly of the genomes was performed using Velvet (6). The genomes were annotated using RAST (7), and scaffolds were mapped on the closest neighbor according to RAST using CONTIGuator (8). Protein annotations were extended using Interproscan (9) and BAGEL3 (10) was used for identification of putative bacteriocin gene clusters.

TABLE 1 Genome features and GenBank accession numbers of the strains

Strain	Species	Source of isolation	Bioproject no.	Accession no.
B4098	<i>Bacillus coagulans</i>	Chinese tomato	PRJNA270593	LQYG00000000
B4100	<i>Bacillus coagulans</i>	Low pH sauce	PRJNA270593	LQYH00000000
B4099	<i>Bacillus coagulans</i>	Indian curry	PRJNA270593	LQYI00000000
B4096	<i>Bacillus coagulans</i>	Tomato supreme	PRJNA270593	LQYJ00000000
B4092	<i>Bacillus licheniformis</i>	Buttermilk powder	PRJNA270588	LQYK00000000
B4090	<i>Bacillus licheniformis</i>	Pea soup	PRJNA270588	LQYL00000000
B4091	<i>Bacillus licheniformis</i>	Mushroom soup	PRJNA270588	LQYM00000000
B4102	<i>Bacillus sporothermodurans</i>	Indian curry	PRJNA270602	LQYN00000000
B4140	<i>Bacillus amyloliquefaciens</i>	Pizza	PRJNA270600	LQYO00000000
B425	<i>Bacillus amyloliquefaciens</i>	Sterilized milk	PRJNA270600	LQYP00000000
B4164	<i>Bacillus licheniformis</i>	Unknown food	PRJNA270588	LQYQ00000000
B4144	<i>Bacillus vallismortis</i>	Quiche	PRJNA270602	LQYR00000000

Nucleotide sequence accession numbers. The genome sequences of the twelve *Bacillus* sp. strains have been deposited as whole-genome shotgun projects at DDBJ/EMBL/GenBank under the accession numbers listed in Table 1.

ACKNOWLEDGMENTS

We thank the NGS sequence facility of the University Medical Center of Groningen (UMCG) for performing the sequencing of the strains.

We thank the Top Institute for Food and Nutrition for contributing to the funding of the project in theme 3: Safety and Preservation.

FUNDING INFORMATION

The Top Institute for Food and Nutrition contributed to the funding of the project in Theme 3: Safety and Preservation.

REFERENCES

- Heyndrickx M. 2011. The importance of endospore-forming bacteria originating from soil for contamination of industrial food processing. *Appl Environ Soil Sci* 11:561975. <http://dx.doi.org/10.1155/2011/561975>.
- Scheldeman P, Herman L, Foster S, Heyndrickx M. 2006. *Bacillus spirothermodurans* and other highly heat-resistant spore formers in milk. *J Appl Microbiol* 101:542–555. <http://dx.doi.org/10.1111/j.1365-2672.2006.02964.x>.
- Eijlander RT, de Jong A, Krawczyk AO, Holsappel S, Kuipers OP. 2014. SporeWeb: an interactive journey through the complete sporulation cycle of *Bacillus subtilis*. *Nucleic Acids Res* 42:D685–D691. <http://dx.doi.org/10.1093/nar/gkt1007>.
- Konz D, Doekel S, Marahiel MA. 1999. Molecular and biochemical characterization of the protein template controlling biosynthesis of the lipopeptide lichenysin. *J Bacteriol* 181:133–140.
- Krawczyk AO, Berendsen EM, Eijlander RT, de Jong A, Wells-Bennik MHJ, Kuipers OP. 2015. Draft genome sequences of four *Bacillus thermoamylovorans* strains isolated from milk and acacia gum, a food ingredient. *Genome Announc* 3(2):e00165-15. <http://dx.doi.org/10.1128/genomeA.00165-15>.
- Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
- Galardini M, Biondi EG, Bazzicalupo M, Mengoni A. 2011. CONTIGuator: a bacterial genomes finishing tool for structural insights on draft genomes. *Source Code Biol Med* 6:11. <http://dx.doi.org/10.1186/1751-0473-6-11>.
- Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A, Nuka G, Pesseat S, Quinn AF, Sangrador-Vegas A, Scheremetjew M, Yong SY, Lopez R, Hunter S. 2014. InterProScan 5: Genome-scale protein function classification. *Bioinformatics* 30:1236–1240. <http://dx.doi.org/10.1093/bioinformatics/btu031>.
- van Heel AJ, de Jong A, Montalbán-López M, Kok J, Kuipers OP. 2013. BAGEL3: automated identification of genes encoding bacteriocins and (non-)bactericidal posttranslationally modified peptides. *Nucleic Acids Res* 41:W448–W453. <http://dx.doi.org/10.1093/nar/gkt391>.